

***In vitro* acaricidal activity of several natural products against ibex-derived *Sarcoptes scabiei*.**

Jesús M. Pérez ^{a,*}, Emiliano N. Jesser ^b, Jorge O. Werdin ^b, Colin Berry ^c, Mohamed A. Gebely ^{c,d}, Raquel Crespo-Ginés ^{a,e}, José E. Granados ^f, Antonio J. López-Montoya ^g

^a Departamento de Biología Animal, Vegetal y Ecología, Universidad de Jaén, Campus Las Lagunillas, s.n., E-23071, Jaén, Spain

^b Departamento de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur, San Juan 670, Bahía Blanca, B 8000CPB, Argentina

^c School of Biosciences, Cardiff University, Museum Avenue, Cardiff, CF10 3AX, UK

^d Department of Parasitology and Animal Diseases, Veterinary Research Institute, National Research Centre, Dokki, Giza 12622, Egypt

^e Instituto de Investigación en Recursos Cinegéticos (IREC-CSIC, UCLM, JCCM), Ronda de Toledo 12, E-13071, Ciudad Real, Spain

^f Centro Administrativo Parque Nacional y Parque Natural Sierra Nevada, Carretera Antigua Sierra Nevada, Km 7, E-18071, Pinos Genil, Granada, Spain

^g Department of Statistics and Operational Research, Jaén University, Campus Las Lagunillas, s.n., E-23071, Jaén, Spain

* author for correspondence: jperez@ujaen.es

ORCID codes

ENJ: 0000-0002-4526-6449; JOW: 0000-0001-5152-3782; CB: 0000-0002-9943-548X;
MAG: 0000-0003-3127-2781; RC-G: 0000-0001-7901-7670; JEG: 0000-0002-9787-
9896; AJLM: 0000-0002-0453-7978; JMP: 0000-0001-9159-0365

ABSTRACT

In this study we analysed the effect of the temperature, diverse strains of *Bacillus thuringiensis*, *Lysinibacillus sphaericus* and nanoformulations with essential plant oils (EONP) on the survival of *Sarcoptes scabiei* mites derived from naturally-infested Iberian ibex (*Capra pyrenaica*). In general, mites maintained at 12°C survived more than those maintained at 35°C (40.7 hr and 31.2 hr, respectively). Mites with no treatment survived 27.6 h on average. Mites treated with *B. thuringiensis* serovar. *konkukian* and geranium EONP showed significant reduction in their survival. Despite the fact that these agents seem to be promising candidates for controlling sarcoptic mange in the field, further research is still needed to get stable, efficient and eco-friendly acaricides.

Keywords: *Bacillus thuringiensis*, essential plant oils, nanoparticles, *Sarcoptes scabiei*, survival

1. Introduction

Sarcoptes scabiei is an astigmatid mite causing a dermal disease, namely sarcoptic mange, in domestic and wild mammalian hosts, including man, worldwide, reaching high morbidity and mortality rates (Bornstein et al., 2001; Pence &

Ueckermann, 2002; Arlian & Morgan, 2017). Transmission of this mite between susceptible hosts may be direct, indirect, or a combination of both (Browne et al., 2022).

Control of this disease in wild populations is a challenging task. Although multiple doses of subcutaneous ivermectin (200-400 µg/kg) is the treatment most commonly used (Rowe et al., 2019), its implementation with free-ranging animals is very difficult from a logistic viewpoint. Moreover, this approach may have undesirable impact on non-target organisms and favours the development of resistance by the mite (Walton et al., 2000), among other “secondary” effects (Moroni et al., 2020).

The development of resistance of *S. scabiei* against acaricidal compounds is increasing (Currie et al., 2004; Mounsey et al., 2010; Andriantsoanirina et al., 2014). Therefore, it is crucial to develop new drugs for treating scabies (Walton et al., 2004). *Bacillus thuringiensis* is a Gram-positive bacterium which produces one or several crystalline proteins referred to as δ-endotoxins (Hill & Pinnock, 1998). After being ingested by susceptible arthropods, the *B. thuringiensis* δ-endotoxin crystals are dissolved in the midgut with a consequent production of activated toxic polypeptides commonly known as δ-endotoxin crystal proteins (Cry proteins) (Höfte & Whiteley, 1989), which may belong to a number of distinct structural families (Crickmore et al., 2021). These toxins seem to disrupt the selective permeability of the cell membrane, which ultimately causes the arthropod death from starvation and/or septicemia (Knowles & Dow, 1993). Like *B. thuringiensis*, *Lysinibacillus sphaericus* is another gram positive bacterium able to produce a range of insecticidal proteins and which exerts its effects in a similar manner (Berry, 2012).

Essential oils (EO) are mixtures of diverse volatile compounds synthesized by plants to protect themselves and are considered as new ecofriendly insecticides, since they may show good biological activity against a number of insect pests, low toxicity to

humans and rapid degradation in the environment (Jesser et al., 2020a). However, EO show some disadvantages such as instability, volatility, and low solubility in water, which may limit their applications (Jesser et al., 2020b).

In recent years, the integration of nanotechnology in the field of biopesticides has garnered significant attention. An innovative method, the nanoformulation of EO, has emerged as a solution to shield active compounds from environmental conditions and prevent the gradual loss of EO (Kumar et al., 2020). Several materials, including proteins, synthetic emulsifiers, polysaccharides (such as starch and chitosan), and polyethers (like polyethylene glycol and poly- ϵ -caprolactone), have been explored for their effectiveness in nanoformulating EO or its constituents (Athanassiou et al., 2018; De Luca, et al., 2021). Polyethylene glycol 6000 (PEG 6000) has been extensively investigated over the past few decades for medical, food industry, and pest control applications. This material showed a broad range of solubility, lacks antigenicity and immunotoxicity, and is easily excreted from living organisms without toxicity concerns. PEG 6000 polymeric nanoparticles loaded with EO (EONP) are considered as one of the most important emerging trends in insect pest control (Campolo et al. 2018; Werdin et al. 2014; 2017).

Nanoformulations of *B. thuringiensis* Cry proteins and EO contribute to increase the activity period of such active compounds in the environment and allow reductions in the amount to be used (De Oliveira et al., 2014; De Oliveira et al., 2021).

The aim of this study was to test the potential acaricide effect of several treatments with *B. thuringiensis* and *Lysinibacillus sphaericus* Cry proteins and essential oils nanoparticles formulations.

2. Materials and methods

Preparation of Bacillus thuringiensis and Lysinibacillus sphaericus

Bacillus thuringiensis and *Lysinibacillus sphaericus* spore/crystal preparations were produced using a method previously described (Jones et al., 2008), by growing the strains in Embrapa medium (Monnerat et al., 2007) until approximately 95% sporulation (judged by phase contrast microscopy) after which spores and crystals were harvested by centrifugation and washed in distilled water before lyophilisation and storage at 4°C. Bacterial strains used were obtained as follows: *Lysinibacillus sphaericus* strain IAB59, *Bacillus thuringiensis* serovar. *higo* strain T44001, *Bacillus thuringiensis* serovar. *israelensis* strain 4Q7, *Bacillus thuringiensis* serovar. *kurstaki* strain HD1 from the *Bacillus* Genetic Stock Center: *Bacillus thuringiensis* GP 138, a kind gift from Prof Alejandra Bravo, UNAM, Mexico; and *Bacillus thuringiensis* serovar. *konkukian* strain 97-27 and *Lysinibacillus sphaericus* strain 2362 from the collection of the Pasteur institute, France.

Preparation of nanoparticles

The EONP was synthesized using the melt-dispersion method, a procedure previously outlined by Werdin González et al. (2014). Initially, 20g of PEG 6000 were heated to 65°C on a hotplate stirrer. Subsequently, 2g of geranium (*Geranium maculatum*) or peppermint (*Mentha piperita*) EO were added to the molten PEG 6000. Geranium and peppermint oils were purchased from Swiss-Just (Switzerland). PEG 6000 was acquired from Merck, Germany. Concurrently, the mixture of PEG 6000 and EO was stirred with a D-500 Handheld Homogenizer (D-lab instrument limited) for 15 minutes at 15,000 rpm. The EONP spontaneously formed when the mixture was cooled to -4°C. After 45 minutes at that temperature, the resulting mixture was thoroughly

ground in a refrigerated mortar box at 0°C, and the product was sifted through a stainless-steel sieve with a mesh size of 230. The EONP were stored in airtight polyethylene pouches at $27 \pm 2^\circ\text{C}$ within desiccators containing calcium chloride for seven days before further experimentation.

EO and EOPN composition

According to Yeguerman et al. (2022), the chemical composition of both pre- and post-formulation essential oils (EOs) was analysed using gas chromatography-mass spectrometry (GC-MS) with an Agilent 7890B gas chromatograph coupled to an Agilent 5977A mass spectrometer. A HP-5MS capillary column (30 m $\times$ 0.25 mm i.d. $\times$ 0.25 μm film thickness) was utilized, with helium serving as the carrier gas at a flow rate of 1.0 mL min⁻¹. The oven temperature was initially set at 50 °C for 2 minutes, then ramped at 5 °C min⁻¹ to 200 °C and held for 15 minutes. Injection block temperature was maintained at 280 °C, and 1 μL aliquots of samples were injected. Ionization energy of 70 eV was used for mass spectrometry, scanning from 35 to 550 m/z. Retention indices (RI) of components were determined using a series of n-alkanes (C8-C20). Further identification was accomplished using NIST 2.0 database. Relative percentages of individual components were calculated by averaging gas chromatography with flame ionization detection (GC-FID) peak areas obtained on a DB-5 column under similar conditions. Key components like α -pinene, limonene, menthol, pulegone, and geraniol were confirmed by comparing with their standard samples (Sigma-Aldrich) via co-injection. Essential oils were extracted from polymeric nanoparticles (EOPN) by dissolving 0.5 g of each sample in 5 mL of distilled water, heating at 65 °C for 30 minutes with magnetic stirring. Upon melting of PEG 6000, 4 mL of petroleum ether was added, and the mixture was stirred for 2 hours. Afterward, the ether phase containing the extracted EOs was collected, diluted to a concentration of

0.001 mg mL⁻¹ (0.1% v/v), and subjected to GC-MS and GC analysis for component identification.

EONP Size Measurement

A Malvern Nano ZS90 instrument was used to determine the size of the EONP. The Polydispersity Index (PDI) was calculated as the square of the standard deviation divided by the square of the mean size, serving as an indicator of the homogeneity or heterogeneity in the size distribution of the particles, following the method described by Pascoli et al. (2018). Each sample, consisting of 0.2 g of EONP, was suspended in 10 mL of distilled water for 30 minutes. Subsequently, the dispersion was filtered using Whatman N° 1 filter paper and allowed to equilibrate for 2 hours. Data were statistically compared using one-way analysis of variance (ANOVA), followed by the LSD test ($N = 4$).

EONP Encapsulation Efficiency

As outlined by Werdin González et al. (2014) the encapsulation efficiency was assessed using spectrophotometric methods. For this, 0.1 g of EONP were individually dissolved in 2 mL of an absolute ethanol-water solution (75:25). The resulting mixture was then centrifuged at 9000 rpm for 10 minutes. The supernatant was carefully collected and subjected to analysis via UV-vis spectrophotometry, employing a Shimadzu UV-1203 photometer with the Kinetics-2-Program Pack (P/N 206-62029-10; Shimadzu Corp., Kyoto, Japan) at a wavelength of 290 nm. This process was repeated for four samples, and the quantity of EO was determined by referring to an appropriate calibration curve for free EO in ethanol.

Encapsulation efficiency (EE) was determined from:

$$EE (\%) = \frac{\text{weight of loaded EO}}{\text{weight of initial EO}} \times 100$$

One way analysis of variance (ANOVA) and LSD were used in order to compare the data (N = 4). The mean physical and chemical characteristics of the EONP formulations are included in tables 2 and 3, respectively.

Mite collection

Ibex (*Capra pyrenaica*) with severe mange in the consolidation and chronic stages (Espinosa et al., 2017), with lesions affecting $\geq 50\%$ of the host skin surface were selected as mite donors (Fig. 1a). The ibex were chemically immobilized with a mixture of ketamine (3 mg/kg) and xylazine (3 mg/kg) (Casas-Díaz et al., 2011), and then euthanized with T-61 Intervet® (combination of embutramide and mebezonium iodide) at a dose of 1ml/1.5 kg.

For mite extraction we painted glass Petri dishes black (14 cm diameter), except in a central circle at the bottom (5.5. cm diameter). Then, a 25 W lamp was placed 7-8 cm below the central circle and several skin pieces from the donor ibex were placed around this circle (Figure 1b). In this way, a temperature gradient was created allowing the mites to concentrate in the central area of the dish (Andrews, 1981) after overnight exposure to the lamp (Fig. 1c). Once the mites left the host skin, the skin pieces were removed (Fig. 1d). The aim of this method was to obtain live mites without manipulating them, therefore, avoiding mechanical damage to mites which could affect their survival.

The first assay (including mite extraction and following counts) was carried out at 12°C and 70% relative humidity (RH). During the remaining assays the plates were maintained in an incubator at 35°C and 45% RH. The number of control plates and treatments of each assay are included in Table 1. After an initial count (including both

live and dead mites), live mites (those showing some kind of movement) were counted twice a day until the death of all the mites.

The acaricidal activity of geranium EONP and peppermint EONP was evaluated at 35 $\mu\text{g cm}^{-2}$ and 70 $\mu\text{g cm}^{-2}$, respectively. The decision to use the concentration of peppermint for this nanoparticle was based on research by Jesser et al. (2020a), which indicated that the bioactivity of geranium EONP was higher than that of peppermint EONP against *Plodia interpunctella* (Lepidoptera: Pyralidae). The nanoparticles were dispersed in the central circle at the bottom base of glass Petri dish.

Statistical analysis

Survival of mites subjected to the different treatments was analysed using the non-parametric Kaplan-Meier estimate via survival curves (Kaplan and Meier, 1958). The Log-rank test (Kleinbaum and Klein, 2012) with Bonferroni correction allowed for multiple pairwise comparisons between the survival curves of each treatment in order to determine significant differences between them.

All statistical analyses were performed using R version 4.3.1 (R Core Team, 2023). We used the `survfit()` function to conduct the Kaplan-Meier estimations with the survival package (Therneau, 2023). The survival curves were drawn using `ggsurvplot()` function of the `survminer` package (Kassambara et al., 2020). The log-rank test was carried out using `survdifff()` function of the survival package. Multiple pairwise comparisons were conducted with the `pairwise_survdifff()` function of the `survminer` package. The statistical significance level set in all statistical analyses was 0.05.

3. Results and discussion

First, the chemical analysis of EOPN revealed that β -citronellol and geraniol were the predominant compounds in geranium EOPN (Table 3). Moreover, components such as linalool, menthone, citronellyl formate, and geranyl formate, which, in the pre-formulation sample, were between 8 and 11%, had a significant reduction after formulation ($<1.7\%$). Additionally, minor components present in the original sample ($<3\%$) were undetectable after formulation. In contrast, menthol emerged as the primary compound in peppermint oil and its nanoparticles, as indicated in Table 3. After formulation, a slight decrease was noted in the concentrations of isomenthone, p-menthen-3-one, and menthol acetate. Furthermore, minor components present in the initial sample ($<6\%$) were not detected after formulation.

Mites in control plates survived, on average, 27.6 h. Mean and median survival times are shown in Table 3. We must take into account that this time measurement started with mite extraction, but, on average, ibex death to laboratory mite extraction took around 10.9 ± 6.1 h. As expected, average survival of mites maintained at low temperature ($12\text{ }^{\circ}\text{C}$) reached the highest values: 40.7 h, compared with that at 35°C : 31.2 hr.

Kaplan-Meier survival analysis is depicted in figures 2 and 3. The overall survival function can be seen in Fig. 2a. Log-rank test (Fig. 2b) showed statistically significant differences in the mean survival time for most of the treatments ($\chi^2 = 38309$, $p < 0.0001$). In particular, *B. thuringiensis* serovar. *konkukian* 97-27 and geranium EONP reduced mite survival significantly. Conversely, mites treated with *B. thuringiensis* GP 138, *B. thuringiensis higo* T44001, Ls IAB59 and peppermint EONP survived more than those maintained in the control plates (Table 4; Fig. 3). The remaining treatments did not show significant differences in mite survival compared with controls.

238

239 4. Discussion

240 The *B. thuringiensis* serovar. *konkukian* and geranium EONP formulation gave
241 promising results in our assays, and could be effective in reducing mite survival time.
242 Different encapsulation strategies (for bacteria, Cry proteins and single spores) aimed to
243 increase ingestion of *B. thuringiensis* Cry proteins, need to be tested. Moreover, host
244 contact time and the effect of the temperature and UV radiation on the persistence of *B.*
245 *thuringiensis* in the field must be addressed before performing *in vivo* assays (de
246 Oliveira et al., 2021). The MXPA patent 02008705 (Ramírez, 2004) is a
247 nanoencapsulation technique of a mixture of *B. thuringiensis* Cry proteins and spores
248 with high residual activity. On the other hand, Ureña-Saborío et al. (2017) performed
249 chitosan/TPP nanoparticles containing bacterial metabolic infiltrates of the strain *B.*
250 *thuringiensis* SER-217, and achieved their efficient release in an aqueous medium,
251 together with increasing protection and stability of such compounds.

252 *B. thuringiensis* strain GP 138, which has previously shown activity against the
253 tick *Rhipicephalus microplus* (Fernández-Ruvalcaba et al., 2010), did not show any
254 reduction in *Sarcoptes scabiei* survival in this study. *B. thuringiensis* *konkukian* strain
255 97-27, however, did show activity. This is a genome-sequenced strain of *B.*
256 *thuringiensis* (Han et al., 2006) that, in contrast to other *B. thuringiensis* strains tested,
257 is not recorded as producing known delta endotoxins or invertebrate-active toxins
258 produced during the vegetative stage of growth. The genome does encode toxins with
259 reported roles in mammalian food poisoning such as CytK, and the tripartite toxins Hbl
260 and Nhe (in common with several strains of *B. thuringiensis*). The activity of these
261 proteins against invertebrates has not been reported and it is possible that these or other,
262 as yet uncharacterised, proteins or small molecule toxins are responsible for its activity

against *S. scabiei* in this study. This possibility warrants further investigation to identify the agent responsible for the activity observed.

This study used geranium and peppermint essential oils (EO) to formulate PEG-6000 nanoparticles, due to their bioactivity against various tick species (An and Tak, 2022; Awad et al., 2022; Klafke et al., 2021; Voronova et al., 2022). In the case of geranium oil, citronellol and geraniol are the important constituents that responsible for the acaricidal activity in *Rhipicephalus annulatus* (Ibrahium, et al., 2022). Similarly, the bioactivity of peppermint EO, attributed to menthol and isomenthone, has been observed against *Tetranychus cinnabarinus* and *Tetranychus urticae* (Abd-Allah et al., 2022). Enan (2001) suggested that the toxicity of constituents of essential oils against insect pests might be related to the octopaminergic nervous system of insects, while de Olivera et al. (1997) proposed that certain monoterpenes inhibit cytochrome P450-dependent monooxygenases. Moreover, Ryan and Byrne (1988) identified a connection between the toxicity of monoterpenes, their capacity to inhibit acetylcholinesterase (AChE), and their effectiveness against insects or ticks.

Regarding the bioactivity of nanoparticles, it was observed that geranium EONP exhibited greater efficacy compared to peppermint EOPN. This result could be attributed to the physicochemical characteristics of the nanoparticles. Peppermint EONP had size of 390 nm and were polydisperse these values are higher than geranium EONP. It is well-known that nanoparticle size plays a crucial role in the penetration of bioactive compounds through the cuticle. Hashem et al. (2018) demonstrated that EO nanoformulations enhance cuticle penetration, allowing products to penetrate insects more easily. Furthermore, the nanoscale size of EONP could extend the exposure time of bioactive compounds to insect pests, covering larger areas of the insect cuticle. Additionally, nanoparticles can alter the delivery pattern of EO active ingredients,

thereby enhancing their efficacy (Iavicoli et al., 2017). Moreover, the encapsulation efficiency (EE) of peppermint EONP was 72%, which is lower than that of geranium EONP. It will probably be necessary to use a higher amount of peppermint EONP for an effective pest management program.

In our study, temperature and relative humidity (RH) were maintained during the different assays. When off the host, *Sarcoptes* mites are unable to use water vapor actively from unsaturated ambient air (Arlian and Veselica, 1979) and, therefore, their survival time is strongly affected by ambient RH (Arlian et al., 1984). Mellanby et al. (1942) found that *S. scabiei* (obtained from human scrapings) did not move when temperature was below 15-16°C, but did so rapidly above 20°C; and heating at 50°C for 10 minutes was enough to exterminate the mite. At cooler temperatures (e.g., 4 °C) black bear-derived mites survived over a week (Niedringhaus et al., 2019). Moreover, at low temperatures, survival of *S. scabiei* increases with relative humidity (Davis & Moon, 1987; Arlian et al., 1989). Thus, environmental conditions (mainly temperature and relative humidity) will strongly affect mite survival when off the host and, therefore, its ability to be transmitted indirectly, to spread, to establish and to persist (Castro et al., 2016; Montecino-Latorre et al., 2019; Loredó et al., 2020; Browne et al., 2021).

5. Conclusion

In conclusion, the acaricidal activity of NP formulations of *B. thuringiensis* *konkukian* strain 97-27, of other *B. thuringiensis* strains and of other plant essential oils at different concentrations deserve to be studied in more detail, before considering *in vivo* assays. In a complementary way, the effect of both temperature and RH on the

survival of ibex-derived mites (when off the host) need to be analysed in depth by maintaining mites *in vitro* at a wider range of such conditions.

Acknowledgements

Authors are indebted to Antonio Rodríguez, José López and Isidro Puga for their help in obtaining biological samples for our study.

Funding

Research activities of JEG and JMP are partially supported by the Junta de Andalucía Government (RNM-118 group of the PAIDI), and by the Jaén University (Action 1b). The AUIP (Asociación Universitaria Iberoamericana de Postgrado) scholarship facilitated EJ's mobility between the National University of the South (Argentina) and the Jaén University (Spain) to conduct this research.

CRediT authorship contribution statement

All the coauthors conceived the work; MAG and CB prepared *Bacillus thuringiensis* and *Lysinibacillus sphaericus* crystal proteins; EJ and JOW prepared nanoparticles containing essential oils, JEG searched and captured donor scabietic ibex; EJ, RC and JMP obtained live mites and performed the assays; EJ, JMP and AJLM analysed the data obtained; all the authors contributed to writing this manuscript.

Declaration of Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships.

Data Availability

Original data are available upon reasonable request.

Ethics approval

Procedures carried out in this work were approved by the regional government (Junta de Andalucía): Project 15/12/2018/163, and also by the Ethics Committee of the Jaén University.

References

- Abd-Allah, G.E., Habashy, M.G., Shalaby, M.M., 2022. Efficacy of mint derivatives, *Mentha spicata* L., against two species of *Tetranychus* spp. (Acari: Tetranychidae) and the predator, *Neoseiulus* sp. Egypt. Acad. J. Biol. Sci. A Entomol., 15, 63–70. DOI: 10.21608/EAJBSA.2022.224349.
- An, H., Tak, J.H., 2022. Miticidal and repellent activity of thirty essential oils and their synergistic interaction with vanillin against *Tetranychus urticae* Koch (Acari: Tetranychidae). Ind. Crops Prod., 182, 114872. <https://doi.org/10.1016/j.indcrop.2022.114872>.
- Andrews, J.R., 1981. The extraction of *Sarcoptes scabiei* from mammalian hosts. J. Parasitol. 67, 753–754.
- Andriantsoanirina, V., Izri, A., Botterel, F., Foulet, F., Chosidow, O., Durand, R., 2014. Molecular survey of knockdown resistance to pyrethroids in human scabies mites. Clin. Microbiol. Infect. 20, O139-O141. <https://doi.org/10.1111/1469-0691.12334>.

356 Arlian, L.G., Morgan, M.S., 2017. A review of *Sarcoptes scabiei*: past, present and
357 future. *Parasites Vectors* 10, 297. DOI: 10.1186/s13071-017-2234-1.

358 Arlian, L.G., Runyan, R.A., Achar, S., Estes, S.A., 1984. Survival and infestivity of
359 *Sarcoptes scabiei* var. *canis* and var. *hominis*. *J. Am. Acad. Dermatol.* 11, 210-
360 215. [https://doi.org/10.1016/S0190-9622\(84\)70151-4](https://doi.org/10.1016/S0190-9622(84)70151-4).

361 Arlian, L.G., Veselica, M.M., 1979. Water balance in insects and mites. *Comp.*
362 *Biochem. Physiol.* 64, 191-200. [https://doi.org/10.1016/0300-9629\(79\)90650-9](https://doi.org/10.1016/0300-9629(79)90650-9).

363 Arlian, L.G., Vyszewski-Moher, D.L., Pole, M.J., 1989. Survival of adults and
364 development stages of *Sarcoptes scabiei* var. *canis* when off the host. *Exp. Appl.*
365 *Acarol.* 6, 181-187. DOI:10.1007/BF01193978.

366 Athanassiou, C.G., Kavallieratos, N.G., Benelli, G., Losic, D., Usha Rani. P., Desneux.
367 N., 2018. Nanoparticles for pest control: Current status and future perspectives. *J.*
368 *Pest Sci.* 91, 1–15. <https://doi.org/10.1007/s10340-017-0898-0>.

369 Awad, S.E., Salah, K.B.H., Jghef, M.M., Alkhaibari, A.M., Shami, A.A., Alghamdi,
370 R.A., Awad, A.E., 2022. Chemical characterization of clove, basil and peppermint
371 essential oils; evaluating their toxicity on the development stages of two-spotted
372 spider mites grown on cucumber leaves. *Life* 12, 1751.
373 <https://doi.org/10.3390/life12111751>.

374 Berry, C., 2012. The bacterium, *Lysinibacillus sphaericus*, as an insect pathogen. *J.*
375 *Invert. Pathol.* 109, 1-10. <https://doi.org/10.1016/j.jip.2011.11.008>.

376 Bornstein, S., Mörner, T., Samuel, W.M., 2001. *Sarcope scabiei* and sarcoptic mange.
377 In: Samuel WM, Pybus MJ, Kocan AA (eds) *Parasitic diseases of wild mammals*,
378 2nd edn. Iowa State University Press, Ames, pp 107-119.

379 Browne, E., Diressen, M.M., Ross, R., Roach, M., Carver, S., 2021. Environmental
380 suitability of bare-nosed wombat burrows for *Sarcoptes scabiei*. Int. J. Parasitol.
381 Parasites Wildl. 16, 37-47. <https://doi.org/10.1016/j.ijppaw.2021.08.003>.

382 Campolo, O., Cherif, A., Ricupero, M., Siscaro, G., Grissa-Lebdi, K., Russo, A., Cucci,
383 L.M., Di Pietro, P., Satriano, C., Desneux, N., Biondi, A., Zappalà, L., Palmeri,
384 V., 2017. Citrus peel essential oil nanoformulations to control the tomato borer,
385 *Tuta absoluta*: Chemical properties and biological activity. Sci. Rep. 7, 13036.
386 <https://doi.org/10.1038/s41598-017-13413-0>.

387 Casas-Díaz, E., Marco, I., López-Olvera, J.R., Mentaberre, G., Lavín, S., 2011.
388 Comparison of xylazine-ketamine and medetomidine-ketamine anaesthesia in the
389 Iberian ibex (*Capra pyrenaica*). Eur. J. Wildl. Res. 57, 887-893. DOI:
390 <https://doi.org/10.1007/s10344-011-0500-7>.

391 Castro, I., de la Fuente, A., Fandos, P., Cano-Manuel, F.J., Granados, J.E., Soriguer,
392 R.C., Alasaad, S., Pérez, J.M., 2017. On the population biology of *Sarcoptes*
393 *scabiei* infesting Iberian ibex (*Capra pyrenaica*). Int. J. Acarol. 42, 7-11.
394 <http://dx.doi.org/10.1080/01647954.2015.1109710>.

395 Crickmore, N., Berry, C., Panneerselvam, S., Mishra, R., Connor, T.R., Bonning, B.C.,
396 2021. A structure-based nomenclature for *Bacillus thuringiensis* and other
397 bacteria-derived pesticidal proteins. J. Invertebr. Pathol. 186, 107438.
398 <https://doi.org/10.1016/j.jip.2020.107438>.

399 Currie, B.J., Harumal, P., McKinnon, M., Walton, S.F., 2004. First documentation of in
400 vivo and in vitro ivermectin resistance in *Sarcoptes scabiei*. Clin. Infect. Dis. 39,
401 e8-e12. <https://doi.org/10.1086/421776>.

402 Davis, D.P., Moon, R.D., 1987. Survival of *Sarcoptes scabiei* (De Geer) stored in three
 403 media at three temperatures. J. Parasitol. 73, 661-662.

404 De Luca, I., Pedram, P., Moeini, A., Cerruti, P., Peluso, G., Di Salle, A., Germann, N.,
 405 2021. Nanotechnology development for formulating essential oils in wound
 406 dressing materials to promote the wound-healing process: a review. Appl. Sci. 11,
 407 1713. <https://doi.org/10.3390/app11041713>.

408 De Oliveira, A.C., Ribeiro-Pinto, L.F., Paumgartten, F.J., 1997. In vitro inhibition of
 409 CYP2B1 monooxygenase by β -myrcene and other monoterpenoid compounds.
 410 Toxicol. Lett., 92, 39-46. DOI: 10.1016/s0378-4274(97)00034-9.

411 de Oliveira, J.L., Campos, E.V.R., Bakshi, M., Abhilash, P.C., Fraceto, L.F., 2014.
 412 Application of nanotechnology for the encapsulation of botanical insecticides for
 413 sustainable agriculture: prospects and promises. Biotechnol. Adv. 32, 1550–1561.
 414 <https://doi.org/10.1016/j.biotechadv.2014.10.010>.

415 de Oliveira, J.L., Fernandes Fraceto, L., Bravo, A., Polanczyk, R.A., 2021.
 416 Encapsulation strategies for *Bacillus thuringiensis*: from now to the future. J.
 417 Agric. Food Chem. 69, 4564–4577. <https://doi.org/10.1021/acs.jafc.0c07118>.

418 Enan, E., 2001. Insecticidal activity of essential oils: octopaminergic sites of action.
 419 Comp. Biochem. Physiol. C. Toxicol. Pharmacol., 130, 325-337.
 420 [https://doi.org/10.1016/S1532-0456\(01\)00255-1](https://doi.org/10.1016/S1532-0456(01)00255-1).

421 Espinosa, J., Ráez-Bravo, A., López-Olvera, J.R., Pérez, J.M., Lavín, S.,
 422 Tvarijonaviciute, A., Cano-Manuel, F.J., Fandos, P., Soriguer, R.C., Granados,
 423 J.E., Romero, D., Velarde, R., 2017. Histopathology, microbiology and the
 424 inflammatory process associated with *Sarcoptes scabiei* infection in the Iberian

425 ibex, *Capra pyrenaica*. Parasites Vectors 10, 596. DOI: 10.1186/s13071-017-
 426 2542-5.

427 Fernández-Ruvalcaba, M., Peña-Chora, G., Romo-Martínez, A., Hernández-Velázquez,
 428 V., Bravo de la Parra, A., Pérez De La Rosa, D., 2010. Evaluation of *Bacillus*
 429 *thuringiensis* pathogenicity for a strain of the tick, *Rhipicephalus microplus*,
 430 resistant to chemical pesticides. J. Insect Sci. 10, 186. DOI:
 431 10.1673/031.010.14146.

432 Han, C.S., Xie, G., Challacombe, J.F., Altherr, M.R., Bhotika, S.S., Bruce, D.,
 433 Campbell, C.S., Campbell, M.L., Chen, J., Chertkov, O., Cleland, C.,
 434 Dimitrijevic, M., Doggett, N.A., Fawcett, J.J., Glavina, T., Goodwin, L.A., Hill,
 435 K.K., Hitchcock, P., Jackson, P.J., Keim, P., Kewalramani, A.R., Longmire, J.,
 436 Malfatti, L.S., McMurtry, K., Meincke, L.J., Misra, M., Moseman, B.L., Mundt,
 437 M., Munk, C., Okinaka, R.T., Parson-Quintana, B., Reilly, L.P., Richardson, P.,
 438 Robinson, D.L., Rubin, E., Saunders, E., Tapia, R., Tesmer, J.G., Thayer, N.,
 439 Thompson, L.S., Tice, H., Ticknor, L.O., Wills, P.L., Brettn, T.S., Gilna, P.,
 440 2006. Pathogenomic sequence analysis of *Bacillus cereus* and *Bacillus*
 441 *thuringiensis* isolates closely related to *Bacillus anthracis*. J. Bacteriol. 188, 3382-
 442 3390. DOI:10.1128/JB.188.9.3382–3390.2006.

443 Hashem, A.S., Awadalla, S.S., Zayed, G.M., Maggi, F., Benelli, G., 2018. *Pimpinella*
 444 *anisum* essential oil nanoemulsions against *Tribolium castaneum* - insecticidal
 445 activity and mode of action. Environ. Sci. Pollut. Res. 25, 18802–18812.
 446 <https://doi.org/10.1007/s11356-018-2068-1>.

447 Hill, C.A., Pinnock, D.E., 1998. Histopathological effects of *Bacillus thuringiensis* on
 448 the alimentary canal of the sheep louse, *Bovicola ovis*. J. Invert. Pathol. 72, 9-20.
 449 DOI: 10.1006/jipa.1998.4761.

450 Höfte, H., Whiteley, H.R., 1989. Insecticidal crystal proteins of *Bacillus thuringiensis*.
 451 Microbiol. Rev. 53, 242–255. DOI: 10.1128/mr.53.2.242-255.1989.

452 Hu, J., Wang, X., Xiao, Z., Bi, W., 2015. Effect of chitosan nanoparticles loaded with
 453 cinnamon essential oil on the quality of chilled pork. LWT - Food Sci. Technol.
 454 63, 519–526. <https://doi.org/10.1016/j.lwt.2015.03.049>.

455 Iavicoli, I., Leso, V., Beezhold, D.H., Shvedova, A.A., 2017. Nanotechnology in
 456 agriculture: Opportunities, toxicological implications, and occupational risks.
 457 Toxicol. Appl. Pharmacol. 329, 96–111.
 458 <https://doi.org/10.1016/j.taap.2017.05.025>.

459 Ibrahim, S.M., Aboelhadid, S.M., Wahba, A.A., Farghali, A.A., Miller, R.J., Abdel-
 460 Baki, A.A.S., Al-Quraishy, S., 2022. Preparation of geranium oil formulations
 461 effective for control of phenotypic resistant cattle tick *Rhipicephalus annulatus*.
 462 Sci. Rep., 12, 11693. <https://doi.org/10.1038/s41598-022-14661-5>.

463 Jesser, E., Yeguerman, C., Stefanazzi, N., Gomez, R., Murray, A.P., Ferrero, A.A.,
 464 Werdin-Gonzalez, J.O., 2020a. Ecofriendly approach for the control of a common
 465 insect pest in the food industry, combining polymeric nanoparticles and post-
 466 application temperature. J. Agric. Food Chem. 68, 5951-5958.
 467 <https://dx.doi.org/10.1021/acs.jafc.9b06604>.

468 Jesser, E., Yeguerman, C., Gili, V., Santillan, G., Murray, A.P., Domini, C., Werdin-
 469 González, J.O., 2020b. Optimization and characterization of essential oil

470 nanoemulsions using ultrasound for new ecofriendly insecticides. ACS Sustain.
 471 Chem. Eng. 8, 7981-7992. <https://dx.doi.org/10.1021/acssuschemeng.0c02224>.
 472 Jones, G.W., Wirth, M.C., Monnerat, R.G., Berry, C., 2008. The Cry48Aa-Cry49Aa
 473 binary toxin from *Bacillus sphaericus* exhibits highly restricted target specificity.
 474 Environ. Microbiol. 10, 2418-2424. [https://doi.org/10.1111/j.1462-](https://doi.org/10.1111/j.1462-2920.2008.01667.x)
 475 2920.2008.01667.x.
 476 Kaplan, E.L., Meier, P., 1958. Nonparametric estimation from incomplete observations.
 477 J. Am. Stat. Assoc. 53, 457-481. <https://doi.org/10.2307/2281868>.
 478 Kassambara, A., Kosinski, M., Biecek, P., 2020. Survminer: drawing survival curves
 479 using 'ggplot2'. R package version 0.4.8. [https://CRAN.R-](https://CRAN.R-project.org/package=survminer)
 480 [project.org/package=survminer](https://CRAN.R-project.org/package=survminer).
 481 Klafke, G.M., Thomas, D.B., Miller, R.J., de León, A.A.P., 2021. Efficacy of a water-
 482 based botanical acaricide formulation applied in portable spray box against the
 483 southern cattle tick, *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae),
 484 infesting cattle. Ticks Tick Borne Dis., 12, 101721. DOI:
 485 10.1016/j.ttbdis.2021.101721.
 486 Kleinbaum, D.G., Klein, M., 2012. Kaplan-Meier survival curves and the Log-Rank
 487 test. In: Survival Analysis. Statistics for Biology and Health. Springer, New York.
 488 https://doi.org/10.1007/978-1-4419-6646-9_2.
 489 Knowles, B.H., Dow, J.A.T., 1993. The crystal d-endotoxins of *Bacillus thuringiensis*:
 490 Models for their mechanism of action in the insect gut. BioEssays 15, 469–476.
 491 DOI: 10.1016/j.toxicon.2006.11.022.

492 Kumar, A., Singh, P., Gupta, V., Prakash, B., 2020. Application of nanotechnology to
 493 boost the functional and preservative properties of essential oils. In: Prakash B
 494 (ed) Functional and preservative properties of phytochemicals. Academic Press,
 495 London, pp 241-267. <https://doi.org/10.1016/B978-0-12-818593-3.00008-7>.

496 Loredó, A.I., Rudd, J.L., Foley, J.E., Clifford, D.L., Cypher, B.L., 2020. Climatic
 497 suitability of San Joaquin kit fox (*Vulpes macrotis mutica*) dens for sarcoptic
 498 mange (*Sarcoptes scabiei*) transmission. J. Wildl. Dis. 56, 126-133. DOI:
 499 10.7589/2019-02-035.

500 Mellanby, K., Johnson, C.G., Bartley, W.C., Brown, P., 1942. Experiments on the
 501 survival and behaviour of the itch mite, *Sarcoptes scabiei* DeG. var. *hominis*.
 502 Bull. Ent. Res. 33, 267-271. <https://doi.org/10.1017/S0007485300026584>.

503 Monnerat, R., Cardoso Batista, A., Telles de Medeiros, P., Soares Martins, É., Melatti,
 504 V.M., Botelho Praça, L., Fiúza Dumas, V., Morinaga, C., Demo, C., Menezes
 505 Gomes, A.C., Falcão, R., Brod Siqueira, C., Oliveira Silva-Werneck, J., Berry, C.,
 506 2007. Screening of Brazilian *Bacillus thuringiensis* isolates active against
 507 *Spodoptera frugiperda*, *Plutella xylostella* and *Anticarsia gemmatilis*. Biol.
 508 Control 41, 291-295. <https://doi.org/10.1016/j.biocontrol.2006.11.008>.

509 Montecino-Latorre, D., Cypher, B.L., Rudd, J.L., Clifford, D.L., Mazet, J.A.K., Foley,
 510 J.E., 2019. Assessing the role of dens in the spread, establishment and persistence
 511 of sarcoptic mange in an endangered canid. Epidemics 27, 28-40.
 512 <https://doi.org/10.1016/j.epidem.2019.01.001>.

513 Moroni, B., Valldeperes, M., Serrano, E., López-Olvera, J.R., Lavín, S., Rossi, L., 2020.
 514 Comment on: “The treatment of sarcoptic mange in wildlife: a systematic review”
 515 Parasites Vectors 13, 471. <https://doi.org/10.1186/s13071-020-04347-0>.

516 Mounsey, K.E., Pasay, C.J., Arlian, L.G., Morgan, M.S., Holt, D.C., Currie, B.J.,
 517 Walton, S.F., McCarthy, J.S., 2010. Increased transcription of glutathione s-
 518 transferases in acaricide exposed scabies mites. *Parasites Vectors* 3, 43.
 519 <https://doi.org/10.1186/1756-3305-3-43>.

520 Niedringhaus, K.D., Brown, J.D., Ternent, M.A., Peltier, S.K., Yabsley, M.J., 2019.
 521 Effects of temperature on the survival of *Sarcoptes scabiei* of black bear (*Ursus*
 522 *americanus*) origin. *Parasitol. Res.* 118, 2767-2772.
 523 <https://doi.org/10.1007/s00436-019-06387-7>.

524 Pascoli, M., Lopes-Oliveira, P.J., Fraceto, L.F., Seabra, A.B., Oliveira, H.C., 2018.
 525 State of the art of polymeric nanoparticles as carrier systems with agricultural
 526 applications: a minireview. *Energ. Ecol. Environ.* 3, 137–148.
 527 <https://doi.org/10.1007/s40974-018-0090-2>.

528 Pence, D.B., Ueckermann, E., 2002. Sarcoptic mange in wildlife. *Rev. Sci. Tech. Off.*
 529 *Int. Epiz.* 21, 385-398.

530 R Core Team, 2023. R: a language and environment for statistical computing. R
 531 Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org/>.

532 Ramirez, L.M., 2004. Technology for encapsulating delta endotoxins of *Bacillus*
 533 *thuringiensis* of the *israelensis* variety for extending the activity thereof on
 534 mosquitoes larvae. MXPA Patent 02008705 A, Dec 6, 2004.

535 Rowe, M.L., Whiteley, P.L., Carver, S., 2019. The treatment of sarcoptic mange in
 536 wildlife: a systematic review. *Parasites Vectors* 12, 99.
 537 <https://doi.org/10.1186/s13071-019-3340-z>.

538 Ryan, M.F., Byrne, O., 1988. Plant-insect coevolution and inhibition of
 539 acetylcholinesterase. *J. Chem. Ecol.*, 14, 1965-1975.
 540 <https://doi.org/10.1007/BF01013489>.

541 Therneau, T., 2023. A package for survival analysis in R. R package version 3.5-5,
 542 <https://CRAN.R-project.org/package=survival>.

543 Ureña-Saborío, H., Madrigal-Carballo, S., Sandoval, J., Vega-Baudrit, J.R., Rodríguez-
 544 Morales, A., 2017. Encapsulation of bacterial metabolic infiltrates isolated from
 545 different *Bacillus* strains in chitosan nanoparticles as potential green chemistry-
 546 based biocontrol agents against *Radopholus similis*. *J. Renew. Mater.* 5, 290–299.
 547 <https://doi.org/10.7569/JRM.2017.634119>.

548 Voronova, N., Horban, V., Bohatkina, V., 2022. The effectiveness of acaricidic drugs
 549 based on herbal raw material. *Ecol. Quest.*, 33, 55-71.
 550 <https://doi.org/10.12775/EQ.2022.003>.

551 Walton, S.F., McKinnon, M., Pizzutto, S., Dougall, A., Williams, E., Currie, B.J., 2004.
 552 Acaricidal activity of *Melaleuca alternifolia* (tea tree) oil. In vitro sensitivity of
 553 *Sarcoptes scabiei* var *hominis* to terpinen-4-ol. *Arch. Dermatol.* 140, 563-566.
 554 DOI: 10.1001/archderm.140.5.563.

555 Walton, S.F., Myerscough, M.R., Currie, B.J., 2000. Studies in vitro on the relative
 556 efficacy of current acaricides for *Sarcoptes scabiei* var. *hominis*. *Trans. R. Soc.*
 557 *Trop. Med. Hyg.* 94, 92-96. DOI: 10.1016/s0035-9203(00)90454-1.

558 Werdin-González, J.O., Jesser, E.N., Yeguerman, C.A., Ferrero, A.A., Fernández Band,
 559 B., 2017. Polymer nanoparticles containing essential oils: New options for
 560 mosquito control. *Environ. Sci. Pollut. Res.* 24, 17006–17015. DOI:
 561 10.1007/s11356-017-9327-4.

562 Werdin-González, J.O., Gutiérrez, M.M., Ferrero, A.A., Fernández Band, B., 2014.
563 Essential oils nanoformulations for stored product pest control: Characterization
564 and biological properties. *Chemosphere* 100, 130–138.
565 <https://doi.org/10.1016/j.chemosphere.2013.11.056>.

566 Yeguerman, C.A., Urrutia, R.I., Jesser, E.N., Massiris, M., Delrieux, C.A., Murray,
567 A.P., Werdin-González, J.O., 2022. Essential oils loaded on polymeric
568 nanoparticles: bioefficacy against economic and medical insect pests and risk
569 evaluation on terrestrial and aquatic non-target organisms. *Envirom. Sci. Pollution*
570 *Res.*, 29, 71412-71426. <https://doi.org/10.1007/s11356-022-20848-0>.

Tables

Table 1. Dates of assays for analyse mite survival and the treatments tested.

Table 2. Average Size (AS), in nanometers, polydispersity (PDI), and Encapsulation Efficiency (EE) of geranium and peppermint EOPN, after 7 Days Post-formulation.

Figures represent mean value $\pm$ standard error.

Table 3. Chemical analysis of pre / post-formulation of the oils from geranium and peppermint.

Table 4. Parameters estimated by treatment via Kaplan-Meier analysis. From left to right: type of treatment, number of mites subjected to this treatment, mean survival time, standard error of the mean and median survival time. Mean and median are measured in hours; se: standard error.

Figure Captions

Figure 1. A: Skin of the scabietic donor ibex. **B:** Several ibex skin pieces were placed into a painted glass Petri dish; note that the center of the plate remains transparent. **C:** the light applied to the bottom of the plate generated a temperature gradient into the plate. This gradient favoured mite migration from the skin to the centre of the plate. **D:** protective wear was needed for skin and plates manipulation.

Figure 2. Kaplan-Meier survival curves. Left graph (A) shows the overall survival curve without considering any treatments. Right graph (B) shows the survival curves by treatment, the p-value of the log-rank test and the pairwise multiple comparisons with Bonferroni correction. Significant differences were indicated by different lowercase letters ($p < 0.05$).

Figure 3. Kaplan-Meier survival curves separated for better visualization.